

Biocillin-150 LA

Amoxicillin 15% injection



*For treatment of gastrointestinal, respiratory
and urinary tract infections caused by
amoxicillin sensitive micro-organisms*


Biocillin-150 LA

Solution for parenteral administration

Amoxicillin is a semisynthetic broad-spectrum penicillin with a bactericidal action against both Gram-positive and Gram-negative bacteria. The range of effect includes *Campylobacter*, *Clostridium*, *Corynebacterium*, *E. coli*, *Erysipelothrix*, *Haemophilus*, *Pasteurella*, *Salmonella*, penicillinase negative *Staphylococcus* and *Streptococcus* spp. The bactericidal action is due to inhibition of cell wall synthesis. Amoxicillin is mainly excreted in urine. A major part can also be excreted in bile.

CONTAINS PER ML

Amoxicillin base 150 mg.



INDICATIONS

Gastrointestinal, respiratory and urinary tract infections caused by amoxicillin sensitive micro-organisms, like *Campylobacter*, *Clostridium*, *Corynebacterium*, *E. coli*, *Erysipelothrix*, *Haemophilus*, *Pasteurella*, *Salmonella*, penicillinase negative *Staphylococcus* and *Streptococcus* spp. in calves, cattle, goats, sheep and swine.

DOSAGE

For intramuscular administration.

General:

1 ml per 10 kg body weight,
repeatable if necessary after 48 hours.

Shake well before use and do not administer more than 20 ml in cattle, more than 10 ml in swine and more than 5 ml in calves, sheep and goats per injection site.

WITHDRAWAL TIMES

- For meat : 21 days.
- For milk : 3 days.

Not for use in sheep and goats producing milk for human consumption.

PACKAGING

Vial of 50 ml and 100 ml.



sales@interchemie.com | interchemie.com

Manufactured by
Interchemie werken "De Adelaar" B.V.
Metaalweg 8, Venray, The Netherlands

XFIA15-EN-01-26082020