

Butachem-50

Buparvaquone 5% injection



For treatment of tick-transmitted theileriosis



Butachem-50

Solution for parenteral administration


Buparvaquone is a second-generation hydroxynaphtaquinone with novel features that make it an effective compound for the therapy and prophylaxis of all forms of theileriosis.

CONTAINS PER ML

Buparvaquone50 mg.

INDICATIONS

For treatment of tick-transmitted theileriosis caused by the intracellular protozoan parasites *Theileria parva* (East Coast fever, Corridor disease, Zimbabwean theileriosis) and *T. annulata* (tropical theileriosis) in cattle. It is active against both the schizont and piroplasm stages of *Theileria* spp. and may be used during the incubation period of the disease, or when clinical signs are apparent.



DOSAGE

For intramuscular administration.

The general dosage is 1 ml per 20 kg body weight.

In severe cases the treatment may be repeated within 48 - 72 hours. Do not administer more than 10 ml per injection site. Successive injections should be administered at different sites.

WITHDRAWAL TIMES

- For meat : 42 days.
- For milk : 2 days.

PACKAGING

Vial of 20 and 50 ml.



sales@interchemie.com | interchemie.com

Manufactured by
Interchemie werken "De Adelaar" B.V.
Metaalweg 8, Venray, The Netherlands

XFIBPS-EN-01-28082020