FLUCONIX-340

Targeted liver fluke control with proven fasciolicidal action

Nitroxinil injection for effective treatment of fascioliasis and blood-feeding nematodes in cattle and sheep, including triclabendazole-resistant flukes





Products and People
you can rely on!

Holland

info@interchemie.com | www.interchemie.com



The main pharmacological action of the active ingredient in Fluconix-340, nitroxinil, is fasciolicidal. The lethal action against *Fasciola hepatica* has been demonstrated *in vitro* and *in vivo* in laboratory animals, and in cattle and sheep. The mechanism of action is due to uncoupling of oxidative phosphorylation. It is also active against triclabendazole-resistant *F. hepatica*.

FLUCONIX-340

Solution for injection



Composition

Contains per ml:

Nitroxinil	340 mg.

Indications

Fluconix-340 is indicated for the treatment of fascioliasis (infestations of mature and immature *Fasciola hepatica*) in cattle and sheep. It is also effective, at the recommended dose rate, against adult and larval infestations of *Haemonchus contortus* in cattle and sheep and *Haemonchus placei*, *Oesophagostomum radiatum* and *Bunostomum phlebotomum* in cattle.

Withdrawal times

- For meat:

	Cattle: 60 days.
	Sheep: 49 days.

Not for use in animals producing milk for human consumption.

Dosage

For subcutaneous administration.

The standard dosage is 10 mg nitroxinil per kg body weight.

Cattle:
1.5 ml of Fluconix-340 per 50 kg of body weight.
Both infected and in-contact animals should be treated, treatment being repeated as considered necessary, though not more frequently than once per month. Dairy cows should be treated at drying off (at least 28 days before calving).

Sheep:
Administer according to the following doses scale:

14 - 20 kg	0.5 ml	41 - 55 kg	1.5 ml
21 - 30 kg	0.75 ml	56 - 75 kg	2.0 ml
31 - 40 kg	1.0 ml	> 75 kg	2.5 ml

In outbreaks of fascioliasis each sheep in the flock should be injected immediately when the presence of the disease is recognised, repeating treatment as necessary throughout the period when infestation is occurring, at intervals of not less than one month.

Packaging

Bottles of 50, 100 and 250 ml.

Products and People
you can rely on!

For full product details and updates, scan the QR code to visit our website

XFINI34-EN-01-01102025