# Imochem-120

The effective antiprotozoal medicine containing imidocarb dipropionate





Highly effective treatment and phrophylaxis of anaplasmosis and babesiosis in cattle, horses, sheep and dogs

# **Imochem-120**


Solution for parental administration

Imidocarb is a diamidine of the carbanalide series of antiprotozoal compounds.

### **CONTAINS PER ML**

### **INDICATIONS**

Imochem-120 is indicated for treatment and prophylaxis of babesiosis in cattle, for treatment of babesiosis and anaplasmosis in sheep, for treatment of babesiosis in horses and dogs and for treatment of anaplasmosis in cattle.



Life cycle of Babesia

# WITHDRAWAL TIMES

- For Meat: 90 days. - For Milk: 21 days (cattle).

Not to be used in sheep producing milk for human consumption

### **PACKAGING**

Vial of 20 and 50 ml.



### **DOSAGE**

For parenteral administration.

### Calves and cattle:

### Babesiosis:

Treatment of babesiosis: 1.0 ml per 100 kg body weight, subcutaneously.

Prevention of babesiosis: 2.5 ml per 100 kg body weight, subcutaneously, one month before exposure.

## Anaplasmosis:

Treatment of anaplasmosis: 2.5 ml per 100 kg body weight, subcutaneously.

Elimination of the carrier state: 4.0 ml per 100 kg body weight, subcutaneously, administered twice with a 14-day interval.

### Horses:

Treatment of Babesia caballi: 2.0 ml per 100 kg body weight, once daily for 2 consecutive days, intramuscularly, Treatment of Babesia equi: 3.5 ml per 100 kg body weight, administered 4 times with a 72-hour interval.

1.0 ml per 100 kg body weight, intramuscularly.

0.5 ml per 10 kg body weight, subcutaneously or intramuscularly. Repeat the dose in 2 weeks, for a total of 2 treatments.













