

Intercam-20

Moloxicam injection



*For treatment of respiratory infections, diarrhoea,
acute mastitis, non-infectious locomotor disorders
and chronic musculo-skeletal disorders*

Intercam-20

Solution for parenteral administration

Meloxicam is a Non-Steroidal Anti-Inflammatory Drug (NSAID) of the oxicam class which acts by inhibition of prostaglandin synthesis, thereby exerting anti-inflammatory, anti-exudative, analgesic and antipyretic effects. It reduces leukocyte infiltration into the inflamed tissue. To a minor extent it also inhibits collagen-induced thrombocyte aggregation. Approximately 50% of an administered dose is eliminated via urine, and the remainder via faeces.

CONTAINS PER ML

Meloxicam 20 mg.

INDICATIONS

Cattle: For use in acute respiratory infection with appropriate antibiotic therapy to reduce clinical signs in cattle. For use in diarrhoea in combination with oral re-hydration therapy to reduce clinical signs in calves of over one week of age and young, non-lactating cattle. For adjunctive therapy in the treatment of acute mastitis, in combination with antibiotic therapy. For the relief of post-operative pain following dehorning in calves.

Pigs: For use in non-infectious locomotor disorders to reduce the symptoms of lameness and inflammation. For adjunctive therapy in the treatment of puerperal septicaemia and toxæmia (mastitis-metritis-agalactia syndrome) with appropriate antibiotic therapy.

Horses: For use in the alleviation of inflammation and relief of pain in both acute and chronic musculo-skeletal disorders. For the relief of pain associated with equine colic.

DOSAGE

Cattle:

Single SC or IV injection at a dosage of 0.5 mg meloxicam/kg body weight (i.e. 2.5 ml/100 kg body weight) in combination with antibiotic therapy or with oral re-hydration therapy, as appropriate.

Pigs:

Single IM injection at a dosage of 0.4 mg meloxicam/kg body weight (i.e. 2.0 ml/100 kg body weight) in combination with antibiotic therapy, as appropriate (can be repeated once after 24 h, if necessary).

Horses:

Single IV injection at a dosage of 0.6 mg meloxicam/kg body weight (i.e. 3.0 ml/100 kg body weight).

WITHDRAWAL TIMES

Cattle: Meat and offal: 15 days;
Milk: 5 days (120 hours)


Pigs: Meat and offal: 5 days.

Horses: Meat and offal: 5 days.

Not authorised to use in horses producing milk for human consumption.

PACKAGING

Vial of 50 ml and 100 ml.



sales@interchemie.com | interchemie.com

Manufactured by
Interchemie werken "De Adelaar" B.V.
Metaalweg 8, Venray, The Netherlands

XFIML2-EN-01-26082020