

Intermectin Super

Ivermectin & Clorsulon solution for injection



*For treatment of round and lung worms,
liver-fluke, eye worms, sucking lice and mange mites*



Intermectin Super

Solution for parenteral administration

IVERMECTIN BELONGS TO THE GROUP OF AVERMECTINS AND ACTS AGAINST ROUNDWORMS AND PARASITES. CLORSULON IS A SULPHONAMIDE WHICH ACTS PRIMARILY AGAINST ADULT AND IMMATURE LIVER FLUKES.

CONTAINS PER ML

Ivermectin 10 mg.
Clorsulon 100 mg.

INDICATIONS

Treatment of gastrointestinal roundworms (adults and fourth-stage larvae), lungworms (adults and fourth-stage larvae), liver fluke (*Fasciola hepatica* and *F. gigantica*; adult stages), eye worms, warbles (parasitic stages), sucking lice and mange mites (scabies) in beef cattle and non-lactating dairy cattle.

DOSAGE

For subcutaneous administration.

General:

1 ml per 50 kg body weight.


WITHDRAWAL TIMES

- For meat : 35 days.

Do not use in non-lactating dairy cows, including pregnant heifers within 60 days of calving.
Do not use in lactating animals producing milk for human consumption.

PACKAGING

Vial of 10 ml, 50 ml, 100 ml, 250 ml and 500 ml.



sales@interchemie.com | interchemie.com

Manufactured by
Interchemie werken "De Adelaar" B.V.
Metaalweg 8, Venray, The Netherlands

XFHCR-EN-01-27082020