

# Limoxin-50

*Oxytetracycline 5% injection*



*For the treatment and control of arthritis,  
gastrointestinal and respiratory infections  
caused by oxytetracycline sensitive micro-organisms*




# Limoxin-50

Solution for parenteral administration

Oxytetracycline belongs to the group of tetracyclines and acts bacteriostatic against many Gram-positive and Gram-negative bacteria like *Bordetella*, *Campylobacter*, *Chlamydia*, *E. coli*, *Haemophilus*, *Mycoplasma*, *Pasteurella*, *Rickettsia*, *Salmonella*, *Staphylococcus* and *Streptococcus* spp. The action of oxytetracycline is based on inhibition of bacterial protein synthesis. Oxytetracycline is mainly excreted in urine, for a small part in bile and in lactating animals in milk.

## CONTAINS PER ML

Oxytetracycline base ..... 50 mg.



## INDICATIONS

Arthritis, gastrointestinal and respiratory infections caused by oxytetracycline sensitive micro-organisms, like *Bordetella*, *Campylobacter*, *Chlamydia*, *E. coli*, *Haemophilus*, *Mycoplasma*, *Pasteurella*, *Rickettsia*, *Salmonella*, *Staphylococcus* and *Streptococcus* spp. in calves, goats, sheep and swine.

## DOSAGE

For intramuscular or subcutaneous administration.

### Full-grown animals:

1 ml per 5 - 10 kg body weight,  
for 3 - 5 days.

### Young animals:

2 ml per 5 - 10 kg body weight,  
for 3 - 5 days.


Do not administer more than 10 ml in swine and more than 5 ml in calves, goats and sheep per injection site.

## WITHDRAWAL TIMES

- For meat : 12 days.
- For milk : 5 days.

## PACKAGING

Vial of 50 and 100 ml.



[sales@interchemie.com](mailto:sales@interchemie.com) | [interchemie.com](http://interchemie.com)



**Manufactured by**  
Interchemie werken "De Adelaar" B.V.  
Metaalweg 8, Venray, The Netherlands

XF105-EN-01-26082020