

Limoxin-100

Oxytetracycline 10% injection



*For the treatment and control of arthritis
gastrointestinal and respiratory infections
caused by oxytetracycline sensitive micro-organisms*


Limoxin-100

Solution for parenteral administration

Oxytetracycline belongs to the group of tetracyclines and acts bacteriostatic against many Gram-positive and Gram-negative bacteria like *Bordetella*, *Campylobacter*, *Chlamydia*, *E. coli*, *Haemophilus*, *Mycoplasma*, *Pasteurella*, *Rickettsia*, *Salmonella*, *Staphylococcus* and *Streptococcus* spp. The action of oxytetracycline is based on inhibition of bacterial protein synthesis. Oxytetracycline is mainly excreted in urine, for a small part in bile and in lactating animals in milk.

CONTAINS PER ML

Oxytetracycline base 100 mg.



INDICATIONS

Arthritis, gastrointestinal and respiratory infections caused by oxytetracycline sensitive micro-organisms, like *Bordetella*, *Campylobacter*, *Chlamydia*, *E. coli*, *Haemophilus*, *Mycoplasma*, *Pasteurella*, *Rickettsia*, *Salmonella*, *Staphylococcus* and *Streptococcus* spp. in calves, cattle, goats, sheep and swine.

DOSAGE

For intramuscular or subcutaneous administration.

Full-grown animals:

1 ml per 10 - 20 kg body weight for 3 - 5 days.

Young animals:

2 ml per 10 - 20 kg body weight for 3 - 5 days.

Do not administer more than 20 ml in cattle, more than 10 ml in swine and more than 5 ml in calves, goats and sheep per injection site.

WITHDRAWAL TIMES

- For meat : 12 days.
- For milk : 5 days.

PACKAGING

Vial of 50 and 100 ml.



sales@interchemie.com | interchemie.com



Manufactured by
Interchemie werken "De Adelaar" B.V.
Metaalweg 8, Venray, The Netherlands

XF1010-EN-01-26082020