

Limoxin-200 LA

Oxytetracycline 20% injection



*For the treatment and control of arthritis
gastrointestinal and respiratory infections
caused by oxytetracycline sensitive micro-organisms*


Limoxin-200 LA

Solution for parenteral administration

Oxytetracycline belongs to the group of tetracyclines and acts bacteriostatic against many Gram-positive and Gram-negative bacteria like *Bordetella*, *Campylobacter*, *Chlamydia*, *E. coli*, *Haemophilus*, *Mycoplasma*, *Pasteurella*, *Rickettsia*, *Salmonella*, *Staphylococcus* and *Streptococcus* spp. The action of oxytetracycline is based on inhibition of bacterial protein synthesis. Oxytetracycline is mainly excreted in urine, for a small part in bile and in lactating animals in milk.

CONTAINS PER ML

Oxytetracycline base 200 mg.



INDICATIONS

Arthritis, gastrointestinal and respiratory infections caused by oxytetracycline sensitive micro-organisms, like *Bordetella*, *Campylobacter*, *Chlamydia*, *E. coli*, *Haemophilus*, *Mycoplasma*, *Pasteurella*, *Rickettsia*, *Salmonella*, *Staphylococcus* and *Streptococcus* spp. in calves, cattle, goats, sheep and swine.

DOSAGE

For intramuscular or subcutaneous administration.

General:

1 ml per 10 kg body weight.

This dosage can be repeated after 48 hours when necessary.

Do not administer more than 20 ml in cattle, more than 10 ml in swine and more than 5 ml in calves, goats and sheep per injection site.

WITHDRAWAL TIMES

- For meat : 28 days.
- For milk : 7 days.

PACKAGING

Vial of 50 and 100 ml.



sales@interchemie.com | interchemie.com



Manufactured by
Interchemie werken "De Adelaar" B.V.
Metaalweg 8, Venray, The Netherlands

XF1020-EN-01-26082020