

Penstrep-400

A combination of penicillin G and dihydrostreptomycin



Arthritis, mastitis and gastrointestinal, respiratory and urinary tract infections caused by penicillin and dihydrostreptomycin sensitive micro-organisms


Penstrep-400

Suspension for parenteral administration

The combination of procaine penicillin G and dihydrostreptomycin acts additive and in some cases synergistic. Procaine penicillin G is a small-spectrum penicillin with a bactericidal action against mainly Gram-positive bacteria like *Clostridium*, *Corynebacterium*, *Erysipelothrix*, *Listeria*, penicillinase-negative *Staphylococcus* and *Streptococcus* spp. Dihydrostreptomycin is an aminoglycoside with a bactericidal action against mainly Gram-negative bacteria like *E. coli*, *Campylobacter*, *Klebsiella*, *Haemophilus*, *Pasteurella* and *Salmonella* spp.

CONTAINS PER ML

Procaine penicillin G 200 000 IU.
Dihydrostreptomycin sulphate 200 mg.



INDICATIONS

Arthritis, mastitis and gastrointestinal, respiratory and urinary tract infections caused by penicillin and dihydrostreptomycin sensitive micro-organisms, like *Campylobacter*, *Clostridium*, *Corynebacterium*, *E. coli*, *Erysipelothrix*, *Haemophilus*, *Klebsiella*, *Listeria*, *Pasteurella*, *Salmonella*, *Staphylococcus* and *Streptococcus* spp. in calves, cattle, goats, sheep and swine.

DOSAGE

For intramuscular administration.

Cattle:

1 ml per 20 kg body weight for 3 days.

Calves, goats, sheep and swine:

1 ml per 10 kg body weight for 3 days.


Note: Shake well before use and do not administer more than 20 ml in cattle, more than 10 ml in swine and more than 5 ml in calves, sheep and goats per injection site.

WITHDRAWAL TIMES

- For kidney : 45 days.
- For meat : 21 days.
- For milk : 3 days.

PACKAGING

Vial of 50 and 100 ml.



sales@interchemie.com | interchemie.com

Manufactured by
Interchemie werken "De Adelaar" B.V.
Metaalweg 8, Venray, The Netherlands

XFIPS-EN-01-26082020