

Penstrep-400 LA

A combination of penicillin G and dihydrostreptomycin



Arthritis, mastitis and gastrointestinal, respiratory and urinary tract infections caused by penicillin and dihydrostreptomycin sensitive micro-organisms


Penstrep-400 LA

Suspension for parenteral administration

The combination of penicillin G and dihydrostreptomycin acts additive and in some cases synergistic. Procaine penicillin G and benzathine penicillin G are small-spectrum penicillins with a bactericidal action against mainly Gram-positive bacteria like *Clostridium*, *Corynebacterium*, *Erysipelothrix*, *Listeria*, penicillinase-negative *Staphylococcus* and *Streptococcus* spp. Dihydrostreptomycin is an aminoglycoside with a bactericidal action against mainly Gram-negative bacteria like *E. coli*, *Campylobacter*, *Klebsiella*, *Haemophilus*, *Pasteurella* and *Salmonella* spp.

CONTAINS PER ML

Procaine penicillin G	100 000 IU.
Benzathine penicillin G	100 000 IU.
Dihydrostreptomycin sulphate	200 mg.



INDICATIONS

Arthritis, mastitis and gastrointestinal, respiratory and urinary tract infections caused by penicillin and dihydrostreptomycin sensitive micro-organisms, like *Campylobacter*, *Clostridium*, *Corynebacterium*, *E. coli*, *Erysipelothrix*, *Haemophilus*, *Klebsiella*, *Listeria*, *Pasteurella*, *Salmonella*, *Staphylococcus* and *Streptococcus* spp.

DOSAGE

For intramuscular administration.

General:

1 ml per 10 kg body weight every 72 hours.

When recommended by professional opinion, it can be administrated every 48 hours.

Note: Shake well before use and do not administer more than 20 ml in cattle, more than 10 ml in swine and more than 5 ml in calves, sheep and goats per injection site.

WITHDRAWAL TIMES

- For kidney : 45 days.
- For meat : 30 days.
- For milk : 5 days.

PACKAGING

Vial of 50 and 100 ml.



sales@interchemie.com | interchemie.com

Manufactured by
Interchemie werken "De Adelaar" B.V.
Metaalweg 8, Venray, The Netherlands
XFIPBS-EN-01-26082020