

# Procaben LA

*Procaine & Benzathine penicillin injection*



*Treatment for arthritis, mastitis and gastrointestinal,  
respiratory and urinary tract infections  
caused by penicillin sensitive micro-organisms*

# Procaben LA

Suspension for parenteral administration

**Procaine and benzathine penicillin G are small-spectrum penicillins with a bactericidal action against Gram-positive and Gram-negative bacteria like *Campylobacter*, *Clostridium*, *Corynebacterium*, *Erysipelothrix*, *Haemophilus*, *Listeria*, *Pasteurella*, penicillinase negative *Staphylococcus* and *Streptococcus* spp. Because of the slow resorption of benzathine penicillin G, the action is maintained for two days.**

## INDICATIONS

Arthritis, mastitis and gastrointestinal, respiratory and urinary tract infections caused by penicillin sensitive micro-organisms, like *Campylobacter*, *Clostridium*, *Corynebacterium*, *Erysipelothrix*, *Haemophilus*, *Listeria*, *Pasteurella*, penicillinase-negative *Staphylococcus* and *Streptococcus* spp. in calves, cattle, goats, sheep and swine.

## CONTAINS PER ML

Procaine penicillin G ..... 150 000 IU.  
Benzathine penicillin G ..... 150 000 IU.

## DOSAGE

For intramuscular administration.

### Cattle:

1 ml per 20 kg body weight.

### Calves, goats, sheep and swine:

1 ml per 10 kg body weight.

This dosage can be repeated after 48 hours when necessary.

Note: Shake well before use and do not administer more than 20 ml in cattle, more than 10 ml in swine and more than 5 ml in calves, sheep and goats per injection site.

## WITHDRAWAL TIMES

- For meat : 14 days.
- For milk : 3 days.

## PACKAGING

Vial of 100 ml.



[sales@interchemie.com](mailto:sales@interchemie.com) | [interchemie.com](http://interchemie.com)



**Manufactured by**  
Interchemie werken "De Adelaar" B.V.  
Metaalweg 8, Venray, The Netherlands

XFIPB-EN-01-27082020